



Occurrence, biology, natural enemies and management of *Tuta absoluta* in Africa

Ramzi Mansour^{1,2}, Thierry Brévault^{3,5#}, Anaïs Chailleux^{4,5#},
Asma Cherif^{6,#}, Kaouthar Grissa-Lebdi^{6,#}, Khalid Haddi^{7.#},
Samira A. Mohamed^{8,#}, Robert S. Nofemela^{9,#}, Abiola Oke^{10,#},
Serigne Sylla^{3,#}, Henri E.Z. Tonnang^{11,#}, Lucia Zappalà^{2,#}, Marc Kenis¹²,
Nicolas Desneux¹³, Antonio Biondi^{2*}

¹ Université de Carthage, Institut Supérieur des Études Préparatoires en Biologie-Géologie (ISEP-BG), Section des Sciences Biologiques, 2036 La Soukra, Tunis, Tunisia

² University of Catania, Department of Agriculture, Food and Environment, 95123 Catania, Italy

³ CIRAD, UPR AIDA, Centre commun ISRA-IRD, Dakar, Senegal

⁴ CIRAD, UPR HortSys, Centre commun ISRA-IRD, Dakar, Senegal

⁵ Université de Montpellier, CIRAD, Montpellier, France

⁶ Université de Carthage, Institut National Agronomique de Tunisie, Laboratoire d'Entomologie-Acarologie, 1082 Cité Mahrajène, Tunis, Tunisia

⁷ Departamento de Entomologia, Universidade Federal de Viçosa, Viçosa, MG 36570-900, Brazil

⁸ International Centre of Insect Physiology and Ecology (ICIPE), PO Box 30772-00100, Nairobi, Kenya

⁹ Agricultural Research Council (ARC) – Plant Protection Research Institute, Private Bag X134, Queenswood 0121, South Africa

¹⁰ National Horticultural Research Institute (NIHORT), PMB 5432, Ibadan, Oyo state, Nigeria

¹¹ International Institute for Tropical Agriculture (IITA), 08 BP 0932-Trip Postal Cotonou, Benin

¹² CABI, 2800 Delémont, Switzerland

¹³ INRA (French National Institute for Agricultural Research), Université Côte d'Azur, CNRS, UMR 1355-7254, Institut Sophia Agrobiotech, Sophia Antipolis, France

These authors contributed equally

* Corresponding author: antonio.biondi@unict.it

With 1 figure and 1 table

Abstract: The South American tomato pinworm, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae), is a devastating invasive pest of tomato crops in several areas around the world including Africa. Here, we comprehensively review and discuss the relevant scientific knowledge on its occurrence, biological and ecological aspects, fortuitous insect natural enemies, and potential advantages and constraints of pest management scenarios adopted against this pest in tomato crops in both Northern and Sub-Saharan Africa. We conclude the manuscript with a comprehensive research agenda providing future priorities towards sustainable control of this important tomato pest.

Keywords: augmentative biological control, classical biological control, conservative biological control, Gelechiidae, invasive pest, Integrated Pest Management, South American tomato pinworm

1 Introduction

Invasive alien insects are generally characterized by successful adaptations to newly invaded areas (Pimentel et al. 2005, Asplen et al. 2015). Some of them are serious threats to agricultural crops, biodiversity and ecosystem functioning, resulting in major economic losses and ecological damages (Pimentel et al. 2000, Sala et al. 2000, Simberloff et al. 2013). Among invasive insects of economic importance, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae), a leafminer native to South America (Guedes & Picanço 2012, Campos et al. 2017, Biondi et al. 2018), is considered a key, devastating pest of processing and fresh tomatoes (*Solanum lycopersicum* L.). This pest was first detected outside South America in 2006 and it is now distributed in most tomato-growing areas in Europe, Africa and Asia (Desneux et al. 2010, 2011, Brévault et al. 2014, Biondi et al. 2018, Han et al. 2018). However, this pest has not been reported in China that is the largest tomato producing country worldwide (Xian et al. 2017, Han et al. 2018).

This Neotropical insect has reached economic pest status in most invaded areas around the world because of its high demographical potential, which is mainly due to a short generation time, a relatively wide host range, a good thermal adaptability and its aptitude to develop insecticide resistance (Desneux et al. 2010, Guedes & Picanço 2012, Tropea Garzia et al. 2012, Biondi et al. 2018). Factors including temperature, predation, parasitism, egg viability, and entomopathogenic agents were proven to affect natural development and survival of this pest (Tonnag et al. 2015, Guimapi et al. 2016, Martins et al. 2016, Bacci et al. 2018). The four larval stages feed on leaves, stems, twigs and fruits, which can cause substantial crop damages and yield losses reaching 100% on tomato crops growing under greenhouse and/or open-field conditions (Desneux et al. 2010, 2011, Biondi et al. 2018). In this perspective, developing and implementing the most appropriate control tactics against this pest is considered a priority for achieving successful and sustainable management of *T. absoluta* worldwide (Campos et al. 2017, Biondi et al. 2018). In Africa, *T. absoluta* has spread

rapidly since its first arrival in three North African countries (Tunisia, Algeria and Morocco), during the 2007–2008 tomato growing season. It is now a well-established major economic pest in both Northern and Sub-Saharan Africa (Desneux et al. 2010, Brévault et al. 2014, Tonnang et al. 2015, Mutamiswa et al. 2017). For this reason and with the aim to find out the most suitable management tactics against this pest, several field surveys and various research activities have been performed in several African countries. Here, we review the most relevant literature on the geographical distribution, the field bio-ecology, the natural enemies and all categories of pest management approaches adopted against *T. absoluta* in Africa. We conclude the manuscript by listing the potential future research priorities with the aim of improving the exploitation of local resources for the sustainable management of this global agroecological and economic problem.

2 Predicting the extent of spread through modeling

In Tonnang et al. (2015), molecular characterization of specimens and geo-reference of occurrence records of *T. absoluta* were combined with biological and ecological data into a CLIMEX model to predict the environmental suitability and potential spread of the pest, with focus to Eastern Africa. Results revealed that factors such as temperature and moisture characterized *T. absoluta* population growth. While, the pest ability to survive the cold, hot, wet and dry stress environment are key factors, which delineate its range. Africa was portrayed to be at high risk of invasion and establishment of the pest. A subsequent study proposed a model to predict the timing of spread of *T. absoluta* invasion across Africa (Guimapi et al. 2016). Using rules based on cellular automata approach, the authors determined, among a number of variables (vegetation, temperature, relative humidity, and area of tomato production), that relative humidity seems to have the strongest influence in enhancing the spread of *T. absoluta* by determining the choice of a location during its invasion and spread (Guimapi et al. 2016). These authors indicated that, in all simulations, when relative humidity is included as parameter, invasion and spread evolution of *T. absoluta* are the closest to the boundaries of natural observations. Simulations by inferring *T. absoluta* natural aptitude to fly long distances demonstrated that the pest was capable of expanding its distribution range to the Republic of South of Africa 10 years after first records in Spain (2006). The projections became validated, as by 2016 *T. absoluta* was reported in the Republic of South Africa (DAFF 2016).

3 Occurrence, ecology and damage

Currently, *T. absoluta* is reported in 41 of the 54 African countries (Fig. 1). However, only in the reports for 22 countries the date of first record of *T. absoluta* was indicated. No further information regarding the year of the first record and/or the



Fig. 1. Occurrence and years of first report of *Tuta absoluta* invasion in Africa, as of August 2018, based on the published literature. Years of first report are indicated as follows: 08 (2008), 09 (2009), 10 (2010), 12 (2012), 13 (2013), 14 (2014), 15 (2015), 16 (2017), 17 (2017).

presence of the pest is available for 32 countries (Fig. 1). The pest probably moved from Northern Africa to Eastern Africa through Sudan, and then spread throughout Sub-Saharan Africa, though this hypothesis is yet to be proved. Below we provide the available information on the spread, host range and field ecology of *T. absoluta* in those countries where the pest has been studied and for which published data are available.

3.1 Northern Africa

Tuta absoluta was first reported in the African continent in three countries of the Maghreb area (Tunisia, Algeria, and Morocco) during the year 2008 before being found in Libya and Egypt in 2009 (EPPO 2008, 2009, 2011, Guenaoui 2008, Moussa et al. 2013, Salama et al. 2015) (Fig. 1). In Tunisia, *T. absoluta* was first detected in October 2008 in open-field tomato crops in the Governorate of Sousse, Center-East region (EPPO 2009, Cherif et al. 2013). Since then, it quickly spread to all tomato-producing areas. Pheromone-based monitoring and sampling of tomato leaves in north-eastern Tunisia during the spring-summer growing season (March – July) showed that *T. absoluta* can have 4 to 5 male flight peaks with higher activity reported in summer during which the monthly temperatures averaged 25–30 °C, while three generations were recorded for eggs and larvae (Cherif & Grissa-Lebdi 2017). However, during the period November – May, *T. absoluta* can have 4 generations on tomatoes under greenhouse conditions of northeastern Tunisia (Grissa-Lebdi et al. 2011). In the same region, three flight peaks of *T. absoluta* males were recorded between January and May in greenhouses, with the highest trap counts recorded in spring (Cherif et al. 2013). In Tunisia, *T. absoluta* can be found on either potato (*Solanum tuberosum* L.), eggplant (*S. melongena* L.) or tomato (mainly), while this pest has a shorter development time on tomato compared to the two other host plants (Cherif et al. 2018d).

In Algeria, *T. absoluta* was first detected in Mostaganem (Northwest) in March 2008 (Guenaoui 2008) before spreading to other tomato-producing areas. In this country, *T. absoluta* can attack a wide range of host plants, in addition to tomato, which are: (i) the Solanaceae (*Solanum elaeagnifolium* Cavanilles, *Solanum nigrum* L., *Datura stramonium* L., *Solanum tuberosum* L., and *Solanum melongena* L.), (ii) the Amaranthaceae (*Chenopodium rubrum* L., *Chenopodium bonus-henricus* L., *Spinacia oleracea* L., and *Beta vulgaris* L.), and (iii) the Fabaceae (*Phaseolus vulgaris* L.) (Drouai et al. 2016). In a tomato greenhouse in northeastern Algeria, it was shown that *T. absoluta* had three generations within a short period, with the first male flight detected in mid-December and the highest densities of all life stages were recorded between mid-March and the end of the crop season, taking into account that during the period December – March, greenhouse temperatures were between 19 and 27 °C (Allache et al. 2012). However, in another greenhouse located in the same region, *T. absoluta* had six generations during the tomato growing season extending from October to May, with the highest numbers of immature stages being reported in mid-April (Allache et al. 2015). Monitoring using pheromone-baited traps showed that the occurrence and densities of *T. absoluta* eggs and larvae on tomato plants depended on both temperature and tomato cultivars grown in greenhouses of Biskra region in northeastern Algeria (Allache et al. 2017).

Tomato crop cultivation constitutes the major horticultural sector in Morocco in terms of planted areas and total production and exports in tons. In this country, *T. absoluta* was first found in the northern coastal (Mediterranean) region of Nador in April 2008, infesting open-field tomatoes (Ouardi et al. 2012). During the growing season 2008–2009, *T. absoluta* has spread rapidly to occur in all Moroccan tomato-producing regions causing damage up to 100% in some processing tomato fields

(Ouardi 2011, Ait Taadaouit et al. 2012, Attrassi 2015). In addition to tomatoes, *T. absoluta* is also infesting potatoes in Morocco (Ouardi 2011). Monitoring based on pheromone traps revealed the presence of the pest year-round at high to very high frequency (Ouardi et al. 2012).

In Egypt, *T. absoluta* was first detected in 2009 in tomatoes in Marsa Matrouh (northwestern Egypt), then the pest rapidly spread to the upper and lower regions of Egypt (Moussa et al. 2013, Salama et al. 2015). In several Egyptian tomato-producing areas, the degree of damage by this insect even reached 100% (Moussa et al. 2013). Under plastic greenhouse conditions in Nasr city (Cairo area, northern Egypt), infestation began in the third week of March and both the highest numbers of *T. absoluta* larvae and percentage of tomato infestation occurred in July (Ata & Megahed 2014). These authors found that *T. absoluta* had four generations on tomato from March to July with mean daily temperatures extended from 19 °C in mid-March to 28 °C in early July. In another study, carried out in open-field tomatoes in Assiut region (Upper Egypt) it was shown that *T. absoluta* can have 13 generations per year (Mohamed 2011).

3.2 Eastern Africa

The spread of *T. absoluta* from Northern to Eastern Africa seems to have transited through Sudan (Fig. 1). In Sudan, the pest was first detected in a tomato greenhouse around Khartoum in 2010, and within less than two years the pest has spread to all tomato-growing regions of North, West and Central Sudan. The pest has also swiftly moved southwards, where it had been reported from southern Kordofan, on the border with the Republic of South Sudan. Fruit damage was reported to range between 80 and 100% in most open-field tomato crops in Khartoum State. In addition to tomato, up to 50% *T. absoluta* damage was observed on potato foliage at Western Umdurman Agricultural Scheme and Karery in Khartoum State (Mohamed et al. 2012). Between 2011 and 2014, a wide survey using pheromone traps was carried out in five regions in Sudan to establish the host plants of *T. absoluta*. The results revealed that the principal host plant was tomato, but also this pest attacked potato, eggplant, Jimson weed (*Datura stramonium* L.) and *S. dubium* Fresen in the family Solanaceae, and broad bean (*Vicia faba* L.) and alfalfa plant (*Medicago sativa* L.) in the family Fabaceae (Mohamed et al. 2015). It is believed that from Sudan, *T. absoluta* migrated to Kenya via Ethiopia.

Tuta absoluta was reported to have traversed into Ethiopia via Sudan in 2012 (Goftishu et al. 2014, Retta & Berhe 2015). Similarly to other countries, widespread surveys were conducted in greenhouse and open-field on tomato crops using pheromone traps and the results confirmed the presence of the pest (Goftishu et al. 2014). According to the same authors, heavy losses in tomato crops were reported in greenhouses in eastern Ethiopia where young tomato plants died before flowering due to heavy infestation by *T. absoluta*. In Kenya, *T. absoluta* was first detected in mid March 2014 in Mpeketoni and Witu, in coastal Kenya, and later in Isiolo and Kirinyaga, in open-field tomatoes by the International Centre of Insect Physiology and Ecology (ICIPE) and by personnel of the Kenya Plant Health Inspectorate Service (KEPHIS),

and officially reported in June of the same year (IPPC 2014). It is believed that the pest expanded its ranges to Kenya through Isiolo that is considered as a gateway town in the Kenya – Ethiopia transit corridor. From Kenya, the pest invaded Tanzania (2014), Uganda (2015) and other countries of the region respectively (Chidege et al. 2016, Tumuhaise et al. 2016, Anonymous 2018).

3.3 Western Africa

In Western Africa, *T. absoluta* was officially first detected in Senegal in 2012 (Pfeiffer et al. 2013, Brévault et al. 2014), then in 2013 in Niger and Cape Verde (Duarte 2013, Adamou et al. 2016), in 2016 in Nigeria and Burkina Faso (EPPO 2016, Borisade et al. 2017, Son et al. 2017), and in 2017 in Ghana (IPPC 2017a) (Fig. 1). However, *T. absoluta* has been thought to be present in most other West African countries including, for example, Benin, Cameroon, Guinea, Ivory Coast, Mali and Togo (USDA 2011, El-Lissy 2014, FAO 2018) (Fig. 1).

In Senegal, a preliminary field survey in the main vegetable-producing area (Niayes) reported that tomato fields were abandoned by farmers during February – May 2013 crop cycle, due to severe damages induced by *T. absoluta* larvae (Brévault et al. 2014). Two years after its first report (Pfeiffer et al. 2013), the pest was reported throughout the country, but mainly on the coastal part where resource availability (almost continuous tomato production for fresh market) and more temperate climate are favorable (Sylla et al. 2017). Adult moths were trapped in some areas in Senegal with very few tomato plants, suggesting that moth populations are extremely mobile or are able to persist on alternative host plants. Larvae were recovered from other solanaceous crops in this country such as potato, eggplant, Ethiopian eggplant (*Solanum aethiopicum* L.), whereas other solanaceous plants, such as pepper and sweet pepper, were consistently less infested (Brévault et al. 2014, Sylla, S., unpublished data). Field monitoring throughout the year in the most infested area in Senegal (Niayes) showed highest incidence of *T. absoluta* populations over the late dry season, from April to June, followed by a significant decline during the rainy season, from July to September (Brévault et al. 2014, Sylla et al. 2017). The decrease of resource availability in the surrounding landscape of monitored fields, especially tomato crops, was the main factor affecting the population dynamics over the rainy season. However, alternative host crops such as eggplant and Ethiopian eggplant can provide reservoirs for residual populations of the pest (Sylla et al. 2018).

Three years after initial detection of *T. absoluta* in Niger, more specifically in Burburkabé and Talkoboye (15 and 70 km from Niamey) a national survey showed that the pest has spread to and established in 7 of the 8 regions of Niger (Adamou et al. 2017). In Nigeria, the largest tomato producer in Sub-Saharan Africa (FAOSTAT 2016), *T. absoluta* was first detected in Daura, Katsina State (northern Nigeria) in April 2015. It was later reported in Kano State in June 2015 and Abeokuta, Ogun State in September 2015 and has since spread to all the other tomato-producing States (Oke et al. 2016, 2017a). *Tuta absoluta* monitoring using water-light traps showed very high pest densities with trap catches up to 4.872 adults per trap per day (Oke et al. 2017a). These authors indicated that such invasion led to a national destruction of

tomato farms, which resulted in the shutdown of Dangote tomato processing industry which was slated to start production in 2015. *Tuta absoluta* caused over 80% loss of tomato production in Nigeria in the first cycle of the season 2016 as consequence of a sporadic attack by this insect in northern parts of the country where the pest has been much restricted (Borisade et al. 2017). Tomato yield losses due to *T. absoluta* invasion and spread in Nigeria is equivalent to 720,000 metric tons, taking into account that the country currently produces 1.8 million tons of tomatoes annually (Sanda et al. 2018). In Cameroon, three regions (Northeast, West and South) were reportedly affected by *T. absoluta* with severe yield losses reaching 100% (FAO 2018).

3.4 Southern and Central Africa

Tuta absoluta is currently established throughout Southern Africa except in Madagascar and Mauritius, while in Central Africa, this pest is present in Democratic Republic of the Congo, Equatorial Guinea, Rwanda and Sao Tome and Principe (FAO 2015, 2018, Luangala et al. 2016, Chidege et al. 2017, IPPC 2017b, 2017c, Mutamiswa et al. 2017, Visser et al. 2017, EPPO 2018) (Fig. 1). *Tuta absoluta* was first recorded in South Africa during August 2016 from synthetic sex pheromone traps in Mpumalanga province set up along the border with Mozambique (DAFF 2016). By May 2017, the pest had invaded open-field and greenhouse tomato crops in all nine provinces of South Africa (DAFF 2017). Following *T. absoluta* invasion, countries that did not have this pest at that time banned tomato (and other Solanaceae crops) imports from South Africa. The impact of this ban on the tomato export market has not been estimated yet. Likewise, the impact of *T. absoluta* on national production of tomatoes is yet to be estimated (Modiselle et al. 2017).

As predicted by Tonnang et al. (2015), *T. absoluta* is a persistent pest in Limpopo and Mpumalanga provinces of South Africa where tomatoes are grown largely under irrigation in open-fields due to favorable weather year-round. Despite the continuous presence of *T. absoluta*, a clear pattern has emerged whereby commercial farmers with access to various classes of registered insecticides are better able to manage *T. absoluta* infestations. The pest is devastating fields of organic and resource-poor smallholder farmers where 50 – 100% crop losses are recorded (Nofemela, R., unpublished data). These losses have a significant impact on household food security, as the profits are used to fund other needs. Likewise, the economic impact of *T. absoluta* on potato production has not been determined in South Africa; all we know at this moment is that high numbers of this pest are recorded in pheromone traps on potato fields and that feeding is limited to foliage (Visser, D., ARC-VOPI, personal communication). In addition, its impact on production of host plants such as pepper and eggplant is still unknown.

In Botswana, *T. absoluta* was first found in a tomato field in the Northeast district in December 2016 (Mutamiswa et al. 2017). In the following year, it was trapped in most regions of the country, both in host crops and wild habitats. Interestingly, *T. absoluta* was detected in areas, such as Moremi Island (Okavango Delta), more than 200 km from the nearest human settlements and agricultural activities (Machekano et al. 2018). In this country, the three wild host plants *Solanum coccineum* (Jacq.),

Solanum supinum (Dunal), and *Solanum aculeatissimum* (Jacq.) were reported, in addition to other abiotic factors (e.g., climate suitability and high thermal tolerance), as major dispersal drivers leading to successful establishment of *T. absoluta* in Botswana (Machekano et al. 2018).

In Zambia, field surveys conducted in 2016 using pheromone delta traps showed that *T. absoluta* is present in 12 districts located in either Northern, Muchinga, Copperbelt, Lusaka, Central or Southern Provinces, with the highest infestation levels reported in the Central province (Luangala et al. 2016). This study showed that some of the surveyed farms in Zambia were highly infested with about 90% of crop damage, and that yields were reduced to zero by the end of the third week of infestation where no insecticide treatments were applied. In 2018, surveys in Zambia showed that, in most cases, farmers had learnt how to control *T. absoluta* on tomato but weekly insecticide treatments were needed (Kenis M., unpublished data). In southwestern Angola (Namibe Province), percent damage of open-field tomato crops induced by *T. absoluta* ranged from 84% to 100% where approximately 256 ha out of 3.500.00 ha under cultivation in 2017 were completely destroyed (Chidege et al. 2017). Knowledge of the damaging potential of this pest in other South and Central African countries is still missing. However, the field scenario could not be much different than the one observed in the other tropical African countries.

4 Fortuitous natural enemies

Since the arrival of *T. absoluta* in Africa, several studies were carried out to assess which indigenous natural enemies are able to attack the invasive pest and could

Table 1. Current knowledge of fortuitous natural enemies associated with, or control options used against, specific life stages of *Tuta absoluta* in Africa based on current bibliography (“x”: associated with / used against; “-”: not associated with / not used against).

Fortuitous natural enemies and IPM tools	Eggs	Larvae	Pupae	Adults
Fortuitous natural enemies				
Parasitoids	x	x	-	-
Predators	x	x	-	-
Natural enemy release				
Parasitoids	x	-	-	-
Predators	x	x	-	-
Pheromone-based tools	-	-	-	x
Prophylactic / cultural tools	-	-	x	x
Insecticide applications				
Biopesticides	-	x	x	-
Synthetic insecticides	-	x	-	-

potentially be used in a biological control approach (Table 1). In this framework, several new host-natural enemies associations have been discovered.

4.1 Northern Africa

A survey of native *Trichogramma* spp. (Hymenoptera: Trichogrammatidae), generalist egg parasitoids, in open-field tomato crops in oases of southwestern Tunisia recorded *Trichogramma bourarachae* Pintureau and Babault parasitizing *T. absoluta* eggs on tomatoes (Zouba et al. 2013b). In another study in Tunisian tomato-growing areas (Center, Centereast and Northeast), two indigenous larval ectoparasitoid species were found developing on *T. absoluta*, *Bracon* sp. (Hymenoptera: Braconidae) attacking mature larvae, and *Necremnus* sp. nr *artyne*s (Hymenoptera: Eulophidae) attacking first, second and third instar larvae, the latter being the most abundant species (Abbes et al. 2014). No egg or pupal parasitoids were recorded in this study. In geothermal tomato greenhouses of southern Tunisia, both adult and nymphal stages of the indigenous predator *Nesidiocoris tenuis* (Reuter) (Hemiptera: Miridae) were found attacking eggs and larvae of *T. absoluta* (Ettaiib et al. 2016). The study demonstrated that this predator (adult and nymphal stages) has a clear preference for *T. absoluta* eggs and that third instar nymphs prefer first and second instar larvae of *T. absoluta* over other leafminer's life stages.

Guenauoui et al. (2011a; 2011b) reported the following species of natural enemies developing or feeding on *T. absoluta* in tomato crops in northwestern Algeria: the eulophid parasitoids *Hemiptarsenus zilahisebessi* Erdos, *N. artyne*s, which was the most abundant parasitoid species, *Neochrysocharis formosa* (Westwood), and *Stenomesus* sp., and the mirid predators *Macrolophus pygmaeus* (Rambur), *M. caliginosus* Wagner, *Dicyphus tamaninii* Wagner, and *N. tenuis*. Furthermore, a survey of natural enemies conducted in tomato greenhouses in the region of Mostaganem (northwestern Algeria) showed the presence of three mirid predators: *N. tenuis*, which was the most abundant species, *M. pygmaeus* and *Dicyphus errans* (Wolff), and the seven following parasitoid species: *N. artyne*s, which was the most frequent and abundant species, *Hyposoter didymator* Thunberg (Hymenoptera: Ichneumonidae), *Neochrysocharis* sp., *Sympiesis* sp., *Diglyphus isaea* Walker (Hymenoptera: Eulophidae), *Bracon* sp., and *Trichogramma* sp. (Boualem et al. 2012). These authors indicated that the second instar larva was the most parasitized life stage, while the lowest parasitism rates were observed in fourth instar larvae.

Studies investigating *T. absoluta* natural enemies in Morocco are scarce. Three predators are frequently used for the biological control of *T. absoluta* in protected tomatoes in this country, i.e., the predatory mirids *N. tenuis* and *M. caliginosus* and the predatory mite *Amblyseius swirskii* Athias-Henriot (Mesostigmata: Phytoseiidae) (ONSSA 2010). However, only *N. tenuis* seems to be a common predator of *T. absoluta* (Elaini 2011, Ouardi et al. 2012).

In Egypt, the most complete study on parasitoids and predators has been that of Eman et al. (2016), who collected the larval parasitoid *Diglyphus* sp., two species of the larval-pupal parasitoid *Elasmus* spp. (Hymenoptera: Eulophidae) and the egg

parasitoid *Telenomus* sp. (Hymenoptera: Platygasteridae). They also abundantly found the predatory bug *N. tenuis* and provided data on its population dynamics and predatory efficiency. Other fortuitous natural enemies of *T. absoluta* found in Egypt include the parasitoids *Bracon nigricans* Szépligeti and *Trichogramma euproctidis* Girault (Zappalà et al. 2013, El-Arnaouty et al. 2014).

4.2 Eastern Africa

Several hymenopterous parasitoid species belonging to different families (Chalcididae, Bethyilidae, Braconidae, Eulophidae) and the mirids *N. tenuis* (Reuter) and *M. pygmaeus* have been found attacking various developmental stages of *T. absoluta* in Kenya, although the parasitism level was low (Kinyanjui & Mohamed, S., unpublished data). The parasitoids *B. nigricans*, *B. hebetor*, *Dolichogenidea appellator* (Telenga) (Hymenoptera: Braconidae), *Ecdamua cademat* (Risbec) (Hymenoptera: Torymidae), *Neochrysocharis formosa* (Westwood), and the predators *N. tenuis* and *Macrolophus* sp. were found associated with *T. absoluta* in Sudan (Mahmoud et al. 2013, Idriss et al. 2018). Moreover, Idriss et al. (2018) evaluated the performance of the two native braconid parasitoids, *B. nigricans* and *D. appellator*, against different immature stages of *T. absoluta* under laboratory conditions. Both parasitoids readily attacked the new host in the laboratory with parasitism levels up to 55% and by showing a differential preference for *T. absoluta* larval instars, i.e., early larval instars for *D. appellator* and later instars for *B. nigricans*. This can support the hypothesis that the *T. absoluta* biological control role of these two parasitoid species is synergistic in the field.

4.3 Western Africa

In Senegal, *N. tenuis* was detected feeding on *T. absoluta* for the first time in 2014 (Sylla et al. 2016). To date, it is the only effective predator detected in Western Africa. Field monitoring in Senegal indicated that 80% of *T. absoluta*-infested tomato fields in the 'Niayes' area also host *N. tenuis*. In Nigeria, the most abundant natural enemy observed in five sampled states was *N. tenuis*, followed by *D. errans* (Oke & Oladigbolu 2018a). A survey of *T. absoluta* natural enemies conducted in Nigeria in September 2015 allowed identifying larval braconid parasitoids of the genera *Apanteles* and *Bracon*, as well as egg parasitoids of the genus *Trichogramma* (Oke et al. 2016).

4.4 Southern and Central Africa

In 2017, surveys for *T. absoluta* natural enemies in heavily infested and abandoned tomato fields in South Africa (Limpopo and Gauteng provinces) recovered six species of indigenous parasitoids and a predatory bug (Nofemela R. unpublished results). No

egg parasitoid was recorded in those surveys. Among the recovered natural enemies, the predator *Rhynocoris segmentarius* (Germar) (Hemiptera: Reduviidae) and the pupal parasitoid *Hockeria* sp. (Hymenoptera: Chalcididae) are being tested for their potential as biological control agents of *T. absoluta* in South Africa.

5 Current integrated management options

In Africa, *T. absoluta* is currently managed mainly by applying synthetic insecticides, but also implementing and integrating other approaches, such as prophylactic and cultural practices, use of pheromone-based trapping systems for early detection, monitoring and/or mass trapping, biological control using either generalist predatory bugs or with egg parasitoids, and application of microbial-based or plant extract-based insecticides (Table 1).

5.1 Prophylaxis and cultural practices

To prevent severe infestations by *T. absoluta*, it is always recommended to integrate prophylactic and cultural control measures in holistic and sustainable Integrated Pest Management (IPM) programs. In Senegal, farmers prefer to avoid growing tomato during late dry season, as it is the riskiest period for *T. absoluta* infestations. Field survey of the pest throughout the year showed a drastic decline of *T. absoluta* populations during the rainy season. This can be due to the decrease of resource availability, especially tomato crops that are less planted during the rainy season owing to susceptibility to foliar diseases. However, alternative hosts, such as eggplant and Ethiopian eggplant, can provide reservoirs for residual populations of the pest. Efforts should be focused on the management of such residual populations, e.g. by removing old eggplant plantations, to slow down the reconstitution of populations at the early dry season (Sylla et al. 2018).

The use of insect-proof screens has been shown to prevent and/or limit *T. absoluta* infestations on host plants in Tunisian (Cherif et al. 2013) and Moroccan tomato greenhouses (Ouardi et al. 2012) as long as nets are well maintained and windows screened. In Nigeria, however, Oke et al. (2016) observed that several commercial screen houses are built with large nets (1 × 4 mm) that allow *T. absoluta* entering into the crop. In Eastern Africa the use of insecticide-impregnated nets has been widely applied against mosquitoes and it was recently introduced to control crop pests, such as aphids (Martin et al. 2013). Although pyrethroid-treated nets have been proven to have sublethal effects on adults of *T. absoluta* (Biondi et al. 2015), this tool has not been implemented in practice yet.

Usually, the use of resistant tomato cultivars in plant breeding programs could be a promising tool for controlling *T. absoluta*, as it was already proven in the tomato variety screening by Sohrabi et al. (2016). In this context, Cherif et al. (2013) suggested the use of tomato cultivars with lower suitability for oviposition by *T. absoluta* (eg.

cv. Shams or cv. Chebli) as a promising prophylactic control tactic against *T. absoluta* in northeastern Tunisian tomato greenhouses. In Morocco, it has been suggested that the destruction of weeds (secondary hosts for *T. absoluta*), crop residues and infested plant material could help avoiding or reduce attacks of greenhouse and open-field tomato crops (Ouardi et al. 2012). Additionally, covering the soil with plastic screens that can help avoiding emergence of adults from buried pupae residing in the soil, and removing infested leaves and secondary shoots are considered as efficient cultural control practices against *T. absoluta* in Tunisia (Abbes et al. 2012).

5.2 Pheromone-based tools: early detection, monitoring and mass trapping

Traps baited with sex pheromone lures can be used for early detection of *T. absoluta* in newly invaded areas (Machekano et al. 2018). When the insect is established in a region, the same tool is then used for pest population monitoring purposes. This involves the follow-up of male flight activity during the growing season, which aims at deciding the most appropriate timing for applying either insecticide treatments or, alternatively, biorational control options (Caparros Megido et al. 2013).

In open-field tomato crops of northeastern Tunisia, *T. absoluta* males are monitored using at least two sex pheromone water traps (Pherodis®) per ha (Cherif & Grissa-Lebdi 2017). These authors indicated that insecticide sprays can be triggered based on pheromone monitoring data: when water trap catches reach a threshold of 50 males per trap per week, insecticides should be applied in tomatoes, as recommended by the Tunisian Ministry of Agriculture. Similarly, Ouardi et al. (2012) suggested, as a component of the national IPM action plan, the use of two pheromone-baited delta traps per ha for monitoring *T. absoluta* in Moroccan tomato crops. In Biskra region (northeastern Algeria), three pheromone delta traps per 400 m² tomato greenhouse (two at the entrances and one in the center) or two pheromone delta traps per 200 m² were used to detect the first adult male flights and to study population fluctuations (Allache et al. 2012, 2017). In Menoufia region (northern Egypt), three water traps baited with *T. absoluta* sex pheromone placed in a 2100 m² tomato field crop area were very useful to follow male flight activity from early April to early July (El-Aassar et al. 2015).

Mass trapping, using 32 traps per ha, was tested against *T. absoluta* in open-field industrial tomatoes in Central Tunisia. Results showed that the pheromone lure TUA-Optima® loaded with 0.8 mg of synthetic pheromone was more attractive to males of *T. absoluta* and should be used with high population levels compared to standard lures Pherodis® and TUA-500® with 0.5 mg of pheromone, which can be used in crops with lower *T. absoluta* population densities (Chermiti & Abbes 2012). According to Ouardi et al. (2012), in Morocco, it is recommended to use 20–25 pheromone traps per ha in tomato greenhouses and 40–50 pheromone traps per ha in open-field tomato crops. In open-field tomato crops of Giza governorate (northern Egypt), the mean percent fruit damage was significantly reduced after applying mass trapping of *T. absoluta* males using 8 pheromone-baited (0.5 mg) water traps per 4200 m² (Taha et al. 2013). Experimental water-light traps had a comparable efficacy in decreasing

the pest damage compared to various combinations of synthetic insecticides used in Nigeria (Oke & Oladigbolu 2018b). However, in South Africa, two pheromone products and one in combination with cypermethrin are registered, but their implementation for mass trapping and/or action threshold is not yet achieved.

A possible alternative to mass trapping is the use of pheromone-based mating disruption technique (i.e. confusing males, disrupting mate location and avoiding/reducing mating). This biorational, residue-free control approach showed great success against *T. absoluta* in tomato greenhouses in both Spain and Italy (Vacas et al. 2011, Cocco et al. 2013). Besides, the performance of attract(lure)-and-kill technique using the combination “synthetic sex pheromone + insecticide” could also be tested as a novel management tool against *T. absoluta* in African tomato-producing areas.

5.3 Releases of natural enemies

Three biological control approaches could potentially be used against *T. absoluta* in Africa (Van Driesche & Bellows 1996). Native natural enemies could be mass-produced for regular releases in field affected by the pest (augmentative biological control). The action of natural enemies present in the crops could be enhanced through the adaptation of specific cultural practices (conservation biological control); and exotic natural enemies from the pest native range could be introduced in Africa for permanent establishment (classical biological control).

Although these biological control options show a promising potential for the control of *T. absoluta* in Africa, so far only augmentative biological control is being applied, and exclusively in Northern Africa. The two categories of natural enemies that are being applied with some success are egg parasitoids and predatory bugs. The egg parasitoids *Trichogramma* spp. were shown to be promising biocontrol agents against *T. absoluta* in Tunisia. A release rate of 20 *T. cacoeciae* per plant was effective in reducing overall insect's life stage densities and infestation levels caused by this pest in protected and open-field tomato crops in the Cap-Bon region, the largest tomato-producing area in Tunisia (Cherif et al. 2018c). In protected tomato crops grown in southwestern Tunisia, a reduction of 87% and 78% in leaf damages was observed after releasing a total of 25,000 adults of either parasitoid *T. cacoeciae* or *T. bourarachae*, respectively (Zouba et al. 2013a). El-Arnaouty et al. (2014) tested the efficiency of three release rates (25, 50 and 75 parasitoids per m²) of the indigenous *T. euproctidis* and *T. achaeae* against *T. absoluta* in northern Egyptian greenhouse tomato crops. They found that both *Trichogramma* species were significantly efficient, particularly at higher release rates, in reducing *T. absoluta* mines, and they suggested including the use of *T. euproctidis*, an autochthonous well-adapted species to local environment, in IPM and biocontrol programs of *T. absoluta* in Egypt.

In addition to egg parasitoids, predatory mirids were shown to be promising biocontrol agents against *T. absoluta* in Northern Africa. Ettaib et al. (2016) demonstrated that releases of *N. tenuis* in heated tomato greenhouses of southwestern Tunisia significantly decreased densities of *T. absoluta* eggs in tomatoes. In Moroccan greenhouses, it is recommended to perform one or two releases of *N. tenuis* at a rate

of 1.5–2 individuals per m² (Ouardi et al. 2012). The use of *N. tenuis* as biocontrol candidate against *T. absoluta* is among the main components of IPM programs in greenhouses of the Souss region in southwestern Morocco (Elaini 2011). However, it should be taken into account that level of predator's efficacy may depend on tomato plant cultivar. Indeed, a recent study showed that predator (*Orius albidipennis* Reuter) efficacy in low dense trichome cultivars was significantly higher than that in high dense trichome cultivars (Salehi et al. 2016).

Interestingly, previous studies provided evidence that the combined use of bio-control agents (i.e. parasitoids, predators and microbials) could enhance their action against *T. absoluta* in tomatoes. Indeed, Kortam et al. (2014) demonstrated that the combined use of the parasitoid *T. achaeae*, the predator *M. caliginosus* and the microbial pesticide *Bacillus thuringiensis* Berliner, at the rates of 25 adults per m², 0.25 adult per m², and 1g per liter, respectively, resulted in a significant reduction of *T. absoluta* mines in greenhouse tomato crops in northern Egypt. Similar studies conducted in northern Egyptian open-field crops showed that the combination “*B. thuringiensis* var. *kurstaki* + release of the parasitoid *T. evanescens* Westwood (70–75 adults/m²) + pheromone mass trapping” significantly decreased densities of *T. absoluta* larvae and related damages in tomatoes (Khidr et al. 2013). Moreover, either combination “biweekly releases of *Trichogramma bactrae* Nagaraja + mass trapping” or “releases of *T. evanescens* + bioinsecticide treatment (Protecto and Spinosad bait)” significantly reduced infestation levels by *T. absoluta* in Egyptian open-field tomatoes (Goda et al. 2015, Rizk 2016).

In Western Africa, so far no *T. absoluta* natural enemies have been marketed, neither mass-reared, for biological control purposes. Nevertheless, some are marketed elsewhere in Africa, such as *N. tenuis* and *Trichogramma* spp. in Northern Africa. Some of the companies have distributors or subsidiaries in Sub-Saharan Africa (Kenya, Mozambique, South Africa, Zambia and Zimbabwe). Therefore, there is a strong potential for the development of *N. tenuis* and/or *Trichogramma* spp. release-based control methods in Western or Southern Africa in the future. Moreover, to the best of our knowledge no exotic natural enemies have been introduced in this area so far.

From a practical point of view, it would be appropriate to use a combination “parasitoids + predators” for strengthening the overall biocontrol action. A previous study in Europe demonstrated that *Trichogramma* parasitoids may not build up populations on the *T. absoluta* tomato system alone, but these parasitoids can be used in combination with the predator *M. pygmaeus* to enhance biological control of this pest in tomatoes (Chailleux et al. 2013). Nonetheless, the possible occurrence of intraguild predation should be taken into account before developing and applying any strategy based on the combined release of natural enemies.

Although several natural enemy species have been found forming new host prey-associations with *T. absoluta* in the new invaded range, including Africa, the search for co-evolved natural enemies of the pest in the native range should also be considered, as currently pursued for several invasive pest species worldwide (Daane et al. 2016, Hardwick et al. 2016, Konopka et al. 2017, Tena et al. 2017, Heimpel & Cock 2018). In this regard, ICIPE (Nairobi, Kenya) and the Federal Ministry for Economic Cooperation and Development (BMZ, Germany), led a *Tuta* IPM funded

project, in partnership with the International Potato Center (CIP), (Lima, Peru). This project has the main aim to import into Kenya the *Microgastrine*, larval parasitoid, *Dolichogenidea gelechiidivoris* (Marsh) (= *Apanteles gelechiidivoris* Marsh, see Salas Gervasio et al. 2019) (Hymenoptera: Braconidae). The initial cohort of the colony of *D. gelechiidivoris*, which is currently maintained at Animal Rearing and Containment Unit (ARCU) of ICIPE, originated from a laboratory colony reared on *T. absoluta* at the CIP entomological laboratory in Lima. Laboratory *D. gelechiidivoris* host stage preference of this parasitoid showed that it prefers the early *T. absoluta* larval instar with an average parasitism rate of over 60% (Mohamed, S., unpublished data). Although field evaluations of this parasitoid are yet to be undertaken, the outstanding laboratory performance of this parasitoid indicates that *D. gelechiidivoris* is a promising classical biological control candidate against *T. absoluta*. Indeed, within its native range this parasitoid species has been recovered with high parasitism rates on *T. absoluta*. For example, in Colombia, *D. gelechiidivoris* is considered the most important biological control agent of *T. absoluta* with field parasitism rates reaching 76.6% (Valencia & Peñaloza 1990, Agudelo & Kaimowitz 1997, Vallejo 1999).

5.4 Insecticide applications and risk of resistance development

The application of insecticide treatments has been the most common control method adopted against *T. absoluta* in African tomato-producing areas. Insecticide formulations that have been used against *T. absoluta* can be based on (i) microbials (bacteria, fungi or virus) and microbial derivatives, (ii) plant extracts (botanicals) or (iii) synthetic chemical active substances, alone or in combinations.

The microbial insecticides *B. thuringiensis* and spinosad exhibited satisfactory control of *T. absoluta* in tomato greenhouses of northeastern Tunisia (Grissa-Lebdi et al. 2011). In semi-natural conditions, *B. thuringiensis* var. *kurstaki* (250 g/hl) caused an average mortality between 72 and 80% on the three first instar larvae of *T. absoluta*, 7 days after treatment, while the new spinosyn, spinetoram (50 cc/hl), caused an average mortality of 85.6% on all *T. absoluta* instar larvae (Hafsi et al. 2012). More recently, Borgi et al. (2016) evaluated the insecticidal potential of a local (Tunisian) strain (P1) of the entomopathogenic fungus *Beauveria bassiana* Balsamo and a spontaneous mutant of this strain (P2) against *T. absoluta* larvae in laboratory conditions. They found that both strains were effective, but P2 showed stronger effect than P1, as it caused 100% of larval mortality within 5-day application. The development of insecticide based on this mutant (P2) should then be recommended as a new tool for the biocontrol of *T. absoluta* in Tunisia (Borgi et al. 2016). Several Tunisian and a Kenyan isolates of *Phthorimaea operculella* granulovirus (PhopGV) showed potential toxicity toward *T. absoluta* larvae in the laboratory, with estimated Lethal Concentration 50% (LC₅₀) ranging from 3.2×10^3 to 1.4×10^5 occlusion body/ml (Ben Tiba et al. 2018).

In Menoufia (northern Egypt) open-field tomato crops, the microbial-based insecticides *B. thuringiensis* (200 g/100 liter) and spinosad (75 ml/100 liter) were the most efficient in terms of control of *T. absoluta* larvae and reduction of related leaf and fruit

damages, compared to the overall control effect of biopesticides based on the entomopathogenic fungi *B. bassiana* and *Metarhizium anisopliae* (Metsch.) (El-Aassar et al. 2015). As such it was proven that the combinations “*B. thuringiensis* var. *kurstaki* + Neem extract”, “*B. thuringiensis* var. *kurstaki* + *Trichoderma harzianum* Rifai”, or “*T. harzianum* + Neem extract” significantly reduced densities and damages of *T. absoluta* larvae in northern Egyptian open-field tomatoes (Khidr et al. 2013).

Extracts of plants grown in Africa may also exhibit insecticidal effects against *T. absoluta*. For example, the extracts of plants collected from the Souss valley in Morocco were tested for their potential in controlling *T. absoluta*. They included *Ricinus communis* L. (Euphorbiaceae), *Argania spinosa* L. (Sapotaceae), *Urtica dioica* L. (Urticaceae), *Thymus vulgaris* L. (Lamiaceae), *Ononis natrix* L. (Fabaceae), *Peganum harmala* L. (Nitrariaceae) and *Lawsonia inermis* L. (Lythraceae). The highest rates of mortality of *T. absoluta* larvae in laboratory conditions were produced by extract of *T. vulgaris* leaves (95% mortality) and the seeds of *R. communis* (58% mortality) (Ait Taadaouit et al. 2012, Nilahyane et al. 2012). Additionally, the essential oil of *Syzygium aromaticum* (L.) buds, a plant grown in Morocco, showed high insecticidal toxicity towards *T. absoluta* larvae in laboratory conditions, hence this plant extract could be incorporated in IPM programs against this pest in Morocco (Benchouikh et al. 2016).

Studies conducted in a Central-East Tunisian tomato greenhouse revealed that the insecticide indoxacarb (50 cc/hl) successfully controlled *T. absoluta* larvae (with more than 95% efficacy nine days after treatment), compared to either triflumuron (50 cc/hl) or diafenthiuron (125 cc/hl) (Braham & Hajji 2012). Besides, in north-eastern Tunisian tomato greenhouses, both chemical pesticides cyromazin (30 g/hl) and flubendiamid (30 g/hl) caused 96 and 77% *T. absoluta* larval mortality, respectively, three weeks after insecticide application, while the efficacy of azadirachtin did not exceed 40% (Cherif et al. 2018a). Among seventeen tested insecticides, the eight active substance formulations chlorantraniliprole, chlorfenapyr, indoxcarb, chlofenapyr mixed with indoxacarb, spinosad, spinosad mixed with abamectin, emamectin benzoate, and imidacloprid provided the most promising control results (> 90% average mortality) against *T. absoluta* larvae occurring in tomato open-field crops in Egypt (Moussa et al. 2013).

Chemical insecticides are mostly used to control *T. absoluta* in Sub-Saharan Africa. The Europe-Africa-Caribbean-Pacific Liaison Committee (COLEACP) initiated the evaluation of several insecticides (COLEACP, 2014, 2015). All countries have different pesticides registered for use against *T. absoluta* in tomatoes, however, in many cases products effective against this pest have yet to be registered. For instance, in 2016 the latest list of the Pesticide Sahelian Committee (CSP) authorized 11 synthetic insecticides in tomatoes: neonicotinoids (acetamiprid and imidacloprid), pyrethroids (lambda-cyhalothrin, cypermethrin and deltamethrin), organophosphates (profenofos, chlorpyrifos-ethyl, chlorpyrifos-methyl), a carbamate (methomyl), and two avermectins (abamectin and emamectin benzoate). Besides, three biopesticides are registered and available in Western Africa for the control of this pest; these are based on *B. thuringiensis*, neem extract and spinosad. A part from the microbial insecticide, most of these insecticides are mostly broad-spectrum for which the real *T. absoluta* control

efficacy in Western Africa has not been described yet. Moreover, some of these insecticides have been banned in Europe for their adverse ecotoxicological effects (Ali et al. 2018), such as profenofos, which is one of the most used chemical active substances in tomatoes in Burkina Faso (Son et al. 2018).

In the Niayes area (northern coast of Senegal), the only method adopted by growers to control tomato insect pests, such as *T. absoluta*, is the application of synthetic (mostly broad-spectrum) insecticides including organophosphates, and to a lesser extent pyrethroids and avermectins. Nevertheless, results indicated that those insecticides were not effective in reducing pest's densities in tomatoes in this area (Diatte et al. 2018). The commercial insecticide on mixture of chlorantraniliprole and lambda-cyhalothrin was the first insecticide recommended and extensively used all over Nigeria for the control of *T. absoluta*. However, control failure was reported soon after. For example, Oke et al. (2017) reported that populations of *T. absoluta* from 7 major tomato-producing areas of Nigeria showed resistance to this agrochemical (resistance ratio ranging from 4.09 to 16.97 times), as well as to lambda-cyhalothrin (resistance ranging from 2.66 to 7.88 times) and to deltamethrin (resistance ranging from 2.23 to 6.24 times). Comparative open-field studies conducted in some Nigerian states (Kano, Kaduna, Katsina, Jigawa and Plateau) revealed that the combination of adult mass trapping, foliar sprays of *B. thuringiensis* and abamectin and applications of *M. anisopliae* to the soil was more efficient in controlling *T. absoluta* than several applications of pyrethroids, chlorantraniliprole, flubendiamide and spirotetramat (Oke & Oladigbolu 2018c). Field surveys conducted in Kirinyaga County (Central Kenya) revealed that 94% of the farmers use synthetic insecticides as main control option for *T. absoluta*, with chlorantraniliprole (200 g/L) being the most applied active substance (Peris et al. 2018). In Central Ethiopian open-field conditions, applications of neem (*Azadirachta indica* (L.)) or *Allium sativum* L. botanical extracts, or the use of the entomopathogen fungus *B. bassiana* caused more than 70% of *T. absoluta* larval mortality, which significantly reduced leaf and fruit infestation (Shiberu & Getu 2018).

Although chemical broad-spectrum insecticide treatments have been commonly applied to cope with *T. absoluta* attacks in tomatoes worldwide (Desneux et al. 2011, Campos et al. 2017, Biondi et al. 2018), this control tactic usually proved to be neither effective nor sustainable, leading to dramatic increases in the number of necessary applications and pest control related costs (Potting et al. 2013) as well as to control failures. Since its arrival in Europe and Northern Africa, insecticide resistance in *T. absoluta* showed to be a major problem and a limiting factor in the development of control systems. Such situation is clearly the result of the selection pressure that this pest faced firstly in South America and subsequently in its areas of recent spread. Indeed, the early cases of resistance were reported from South America as a consequence of the intensive use, and overuse of organophosphate and pyrethroid insecticides (Salazar & Araya 1997, 2001, Siqueira et al. 2000, Lietti et al. 2005). Successively, low to moderate levels of indoxacarb resistance as well as high resistance to chitin synthesis inhibitors were detected (Silva et al. 2016). Later on, spinosad resistance reached high levels (Reyes et al. 2012, Campos et al. 2014) and more recently, resistance to diamides is spreading (Campos et al. 2015, Roditakis et al. 2018).

This scenario, where changes in the patterns of insecticide resistance followed the introduction of new chemistries to be used against *T. absoluta*, seems to be occurring in the newly invaded areas. In fact, at the time of its initial establishment in South European and North African countries, the control of *T. absoluta* was based mainly on broad-spectrum insecticides such as pyrethroids due to lack of specifically registered molecules. Thereafter, field collected populations in those regions showed high levels of resistance to pyrethroids and organophosphates (Haddi et al. 2012, 2017, Roditakis et al. 2013a) though these phenomena are more likely to be existent pre-invasion and introduced with the arrival of the first individuals in the area. Furthermore, recent reports of control failure to oxidiazines (Roditakis et al. 2013b), spinosyns and diamides (Roditakis et al. 2015, 2018) are clearly illustrating the high potential of this pest to develop resistance to multiple classes of insecticides.

In the Sub-Saharan countries, although only very few data on the cases of insecticide resistance are currently available, it is highly likely that the invasive nature and destructive potential of this pest, combined with the over-reliance on chemical control will lead to similar outcomes. In addition to resistance, other major pitfall linked to insecticide applications is that these chemical substances show detrimental side effects on native natural enemies, hampering their beneficial effects on pest populations and allowing the emergence of secondary pests (Arnò & Gabarra 2011, Biondi et al. 2012, 2013b, Wanumen et al. 2016, Cherif et al. 2018b, and see Desneux et al. 2007). Furthermore, insecticides can also exhibit adverse side effects on pollinators of tomato, including bumblebees and stingless bees (Mommaerts et al. 2006, Sarto et al. 2014, Ceuppens et al. 2015).

6 Research agenda for sustainable management

The invasive South American tomato pinworm is arguably the key, most devastating insect pest occurring in tomato-producing areas in Northern and Sub-Saharan Africa. In these geographical areas, *T. absoluta* has continued to spread rapidly since its first arrival in 2008, causing serious damages, substantial economic losses and, in certain circumstances, important food security issues. Although this alarming phytosanitary situation has incited researchers, growers and policy makers to deploy great efforts in an attempt to find the best approaches and tactics for suitably managing this major invasive pest, many crucial studies and actions are strongly warranted. For instance, the pest status in Western and Central African countries is still controversial and needs to be clarified as soon as possible. Moreover, the invasion history in this continent, as well as the population genetic structure behind this invasion is still unknown. Specific studies are thus needed in order to clarify these aspects. Untangling the genetic structure of the invading *T. absoluta* population(s) in Africa would also help understanding the propensity of this pest to develop insecticide resistance. This aspect is crucial because insecticide applications are still the most common approach adopted by growers to control *T. absoluta* in Africa. In South America and Europe, the extensive use of synthetic insecticides has led to the development

of resistance to most insecticide classes and, in this regard, Africa is likely to face a similar outcome if nothing is done to reduce the overdependence on synthetic pesticides. This inevitably leads to higher crop protection costs and to side effects on non-target organisms, including farmers and consumers. For these reasons, the use of reduced-risk insecticide compounds, such as plant extracts or microbial-based selective insecticides (especially when locally available) and/or biorational alternatives are the most recommended approaches for achieving an effective control of *T. absoluta* in Africa and worldwide. Among the alternative control measures, the correct use of insect-proof nets (impregnated with insecticides or not), the use of pheromone-based tools (especially for mass trapping), the conservation and augmentation of indigenous biological control agents (especially generalist predators and egg parasitoids) should be integrated in *T. absoluta* IPM packages.

However, much more efforts are needed from researchers, policy makers, IPM practitioners, non-profit organizations and private companies to optimize the implementation of these tools. For instance, future actions should target: the development of standard protocols and the implementation of mass production of locally available biopesticides, based on plant extracts and/or microbials, and natural enemies (generalist predators and/or egg parasitoids); the identification of the optimal trap density for mass trapping; the potential identification of specific habitat management strategies (within farm and at the landscape level) in a conservation biological control approach aiming at increasing the natural presence of parasitoids and predators; further pre-release and post-release testing on the imported South American parasitoid, *D. gelechiidivoris*, and, in case of successful establishment in the new environment, satisfactory *T. absoluta* biological control, and least non-target effects, its intentional diffusion in other tomato-growing African areas should be promoted; the eventual identification of further species-specific and effective natural enemy species in the native range of *T. absoluta*, which could be considered for imports in Africa. Finally, the use of existing non-chemical control strategies needs to be promoted. The awareness of farmers and the governments should be significantly raised on the economic and environmental risks linked with the extensive use of pesticides and on the efficacy and sustainability of those alternative control tools. This would ultimately lead to economically profitable and sustainable control systems, thus increasing the quality of the life of small farmers and of their families, the local consumers as well as the quality of the tomatoes grown for extra-Africa exports.

Acknowledgments: AB and LZ obtained funding from the University of Catania (Project 2016–18 “Emergent Pests and Pathogens and Relative Sustainable Strategies – 5A722192113), from the Italian Ministry of Education, University and Research (MIUR) (PRIN project 2015 “BIOPIC”, 2015BABFCF). AB, LZ and ND thank the ERA-NET action ARIMNET2 2015 call (project “Sustainable TomatoProduction” – STomP). MK was financially supported by the Department for International Development (DFID), UK and the Directorate-General for International Cooperation (DGIS), Netherlands through CABI’s Action on Invasives programme. KH was supported by CAPES Foundation, the Brazilian National Council of Scientific and Technological Development (CNPq). SAM was financially support by the Federal Ministry for Economic Cooperation and Development (BMZ, Germany) (Project No.: 12.1433.7-001.00) through the International Centre of Insect Physiology and Ecology (ICIPE).

References

- Abbes, K., Biondi, A., Zappalà, L. & Chermiti, B. (2014): Fortuitous parasitoids of the invasive tomato leafminer *Tuta absoluta* in Tunisia. – *Phytoparasitica* 42: 85–92.
- Abbes, K., Harbi, A. & Chermiti, B. (2012): The tomato leafminer *Tuta absoluta* (Meyrick) in Tunisia: current status and management strategies. – *EPPO Bulletin* 42: 226–233.
- Adamou, H., Adamou, B., Garba, M., Oumarou, S., Gougari, B., Abou, M., Kimba, A. & Delmas, P. (2016): Confirmation of the presence of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Niger (West Africa). – *International Journal of Environmental Science and Technology* 5: 4481–4486.
- Adamou, H., Garba, M., Dan, Mairo M., Adamou, B., Oumarou, S., Gougari, B., Kimba, A., Abou, M. & Delmas, P. (2017): Geographical distribution of the tomato borer, *Tuta absoluta* Meyrick (Lepidoptera: Gelichiidae) in Niger. – *Scholars Academic Journal of Biosciences* 5: 108–113.
- Agudelo, L. & Kaimowitz, D. (1997): Tecnología agrícola sostenible: Restos institucionales y metodológicos. Dos estudios de Caso en Colombia. – Instituto Interamericano de Cooperación para la Agricultura IICA. San José, Costa Rica.
- Ait, Taadaouit N., Hsaine, M., Rochdi, A., Nilahyane, A. & Bouharroud, R. (2012): Effet des extraits végétaux méthanoliques de certaines plantes marocaines sur *Tuta absoluta* (Lepidoptera, Gelechiidae). – *EPPO Bulletin* 42: 275–280.
- Ali, D., Ali, H., Alifiri, S., Alkahtani, S., Alkahtane, A.A. & Huasain, S.A. (2018): Detection of oxidative stress and DNA damage in freshwater snail *Lymnaea leuteola* exposed to profenofos. – *Frontiers of Environmental Science & Engineering* 12: 1.
- Allache, F., Houhou, M.A., Osmane, I., Naili, L. & Demnati, F. (2012): Suivi de l'évolution de la population de *Tuta absoluta* Meyrick (Gelichiidae), un nouveau ravageur de la tomate sous serre à Biskra (sud-est d'Algérie). – *Faunistic Entomology* 65: 149–155.
- Allache, F., Bouta, Y. & Demnati, F. (2015): Population development of the tomato moth *Tuta absoluta* (Lepidoptera: Gelechiidae) in greenhouse tomato in Biskra, Algeria. – *Journal of Crop Protection* 4: 509–517.
- Allache, F., Demnati, F. & Houhou, M.A. (2017): Population changes of *Tuta absoluta* (Gelechiidae) and fruit loss estimates on three tomato cultivars in greenhouses in Biskra, Algeria. – *Environmental and Experimental Biology* 15: 201–208.
- Anonymous (2018): Somaliland farmers distressed by insects destroying crops in Salahley. An online article, Radio Ergo, published on Monday July 16th, 2018. https://www.hiiraan.com/news4/2018/July/159104/somaliland_farmers_distressed_by_insects_destroying_crops_in_salahley.aspx Accessed on 22 July 2018
- Arnò, J. & Gabarra, R. (2011): Side effects of selected insecticides on the *Tuta absoluta* (Lepidoptera: Gelechiidae) predators *Macrolophus pygmaeus* and *Nesidiocoris tenuis* (Hemiptera: Miridae). – *Journal of Pest Science* 84: 513–520.
- Asplen, M.K., Anfora, G., Biondi, A., Choi, D.-S., Chu, D. et al. (2015): Invasion biology of spotted wing *Drosophila* (*Drosophila suzukii*): a global perspective and future priorities. – *Journal of Pest Science* 88: 469–494.
- Ata, T.E. & Megahed, M.M.M. (2014): Population density of tomato leafminer *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) under protected cultivation in Egypt. – *Middle East Journal of Agriculture Research* 3: 1242–1247.
- Attrassi, K. (2015): Study of the evaluation of damage caused by *Tuta Absoluta* (Meyrick) (Lepidoptera: Gelichidae) on tomato field in Morocco. – *International Journal of Emerging Technologies in Learning* 2: 2046–2053.
- Bacci, L., Silva, E., Martins, J., Soares, M., Campos, M. & Picanço, M. (2018): Seasonal variation of natural mortality factors of *Tuta absoluta* (Lepidoptera: Gelechiidae) in open field tomato cultivation. – *Journal of Applied Entomology*, doi: 10.1111/jen.12567
- Benchouikh, A., Allam, T., Kribii, A., Ounine, K. (2016): L'étude de l'effet insecticide de l'huile essentielle de *Syzygium aromaticum* L. contre les larves de *Tuta absoluta*. – *International Journal of Innovative Research in Science* 20: 188–194.

- Ben Tiba, S., Larem, A., Laarif, A., Fritsch, E., Undorf-Spahn, K., Mohamed, S.A., Ekesi, S., Wennmann, J.T., Kroschel, J., Fattouch, S. & Jehle, J.A. (2018): The potential of novel African isolates of *Phthorimaea operculella* granulovirus for the control of *Tuta absoluta*. – Journal of Applied Entomology, doi: 10.1111/jen.12568
- Biondi, A., Desneux, N., Siscaro, G. & Zappalà, L. (2012): Using organic-certified rather than synthetic pesticides may not be safer for biological control agents: selectivity and side effects of 14 pesticides on the predator *Orius laevigatus*. – Chemosphere 87: 803–812.
- Biondi, A., Desneux, N., Amiens-Desneux, E., Siscaro, G. & Zappalà, L. (2013a): Biology and developmental strategies of the Palaearctic parasitoid *Bracon nigricans* (Hymenoptera: Braconidae) on the neotropical moth *Tuta absoluta* (Lepidoptera: Gelechiidae). – Journal of Economic Entomology 106: 1638–47.
- Biondi, A., Zappalà, L., Stark, J.D. & Desneux, N. (2013b): Do biopesticides affect the demographic traits of a parasitoid wasp and its biocontrol services through sublethal effects? – PLoS One 8: e76548.
- Biondi, A., Zappalà, L., Desneux, N. et al. (2015): Potential toxicity of α -cypermethrin-treated nets on *Tuta absoluta* (Lepidoptera: Gelechiidae). – Journal of Economic Entomology 108: 1191–1197.
- Biondi, A., Guedes, R.N.C., Wan, F.H. & Desneux, N. (2018): Ecology, worldwide spread, and management of the invasive South American tomato pinworm, *Tuta absoluta*: past, present, and future. – Annual Review of Entomology 63: 239–258.
- Borisade, O.A., Kolawole, A.O., Adebo, G.M. & Uwaidem, Y.I. (2017): The tomato leafminer (*Tuta absoluta*) (Lepidoptera: Gelechiidae) attack in Nigeria: effect of climate change on over-sighted pest or agro-bioterrorism? – Journal of Agricultural Extension and Rural Development 9: 163–171.
- Borgi, I., Dupuy, J.W., Blibech, I., Lapailierie, D., Lomenech, A.M., Rebai, A., Ksantini, M., Bonneau, M. & Gargouri, A. (2016): Hyper-proteolytic mutant of *Beauveria bassiana*, a new biological control agent against the tomato borer. – Journal of Agricultural Extension and Rural Development 36: 60.
- Boualem, M., Allaoui, H., Hamadi, R. & Medjahed, M. (2012): Biologie et complexe des ennemis naturels de *Tuta absoluta* à Mostaganem (Algérie). – EPPO Bulletin 42: 268–274.
- Braham, M. & Hajji, L. (2012): Management of *Tuta absoluta* (Lepidoptera, Gelechiidae) with Insecticides on Tomatoes. Insecticides. – Pest Engineering, Farzana Perveen (ed.), ISBN: 978–953–307-895–3, InTech, Croatia, pp. 1–23.
- Brévault, T., Sylla, S., Diatte, M., Bernadas, G. & Diarra, K. (2014): *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae): a new threat to tomato production in sub-Saharan Africa. – African Entomology 22: 441–444.
- Campos, M.R., Rodrigues, A.R., Silva, W.M., Silva, T.B., Silva, V.R. et al. (2014): Spinosad and the tomato borer *Tuta absoluta*: a bioinsecticide, an invasive pest threat, and high insecticide resistance. – PLoS One 9: e103235.
- Campos, M.R., Silva, T., Silva, W.M., Silva, J.E. & Siqueira, H.A. (2015): Susceptibility of *Tuta absoluta* (Lepidoptera: Gelechiidae) Brazilian populations to ryanodine receptor modulators. – Pest Management Science 71: 537–544.
- Campos, M.R., Biondi, A., Adiga, A., Guedes, R.N.C. & Desneux, N. (2017): From the Western Palaearctic region to beyond: *Tuta absoluta* ten years after invading Europe. – Journal of Pest Science 90: 787–796.
- Caparros Megido, R., Haubruge, E. & Verheggen, F.J. (2013): Pheromone-based management strategies to control the tomato leafminer, *Tuta absoluta* (Lepidoptera: Gelechiidae). A review. – Biotechnology, Agronomy, Society and Environment 17: 475–82.
- Ceuppens, B., Eraerfs, M., Vleugels, T., Cnops, G., Roldan-Ruiz, I. & Smagghe, G. (2015): Effects of dietary lambda-cyhalothrin exposure on bumblebee survival, reproduction, and foraging behavior in laboratory and greenhouse. – Journal of Pest Science 88: 777–783.
- Chaillieux, A., Desneux, N., Seguret, J., Khanh, H.Do.T., Maignet, P. & Tabone, E. (2012): Assessing European egg parasitoids as a mean of controlling the invasive South American tomato pinworm *Tuta absoluta*. – PLoS One 7: e48068.

- Chailleux, A., Biondi, A., Han, P., Tabone, E. & Desneux, N. (2013): Suitability of the pest-plant system *Tuta absoluta* (Lepidoptera: Gelechiidae)-tomato for *Trichogramma* (Hymenoptera: Trichogrammatidae) parasitoids and insights for biological control. – Journal of Economic Entomology 106: 2310–2321.
- Cherif, A., Mansour, R. & Grissa-Lebdi, K. (2013): Biological aspects of tomato leafminer *Tuta absoluta* (Lepidoptera: Gelechiidae) in conditions of Northeastern Tunisia: possible implications for pest management. – Environmental and Experimental Biology 11: 179–184.
- Cherif, A. & Grissa-Lebdi, K. (2017): Population dynamics of the tomato leaf miner *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Tunisia natural conditions. – Journal of entomology and zoology studies 5: 427–432.
- Cherif, A., Harbaoui, K., Zappalà, L. & Grissa-Lebdi, K. (2018a): Efficacy of mass trapping and insecticides to control *Tuta absoluta* in Tunisia. – Journal of Plant Diseases and Protection 125: 51–61.
- Cherif, A., Ichaoui, O., Barhoumi-Attia, S. & Grissa-Lebdi, K. (2018b): Life-stage-dependent side effects of selected insecticides on *Trichogramma cacoeciae* (Marchal) (hymenoptera: Trichogrammatidae) under laboratory conditions. – Phytoparasitica 46: 105–113.
- Cherif, A., Mansour, R., Barhoumi-Attia, S., Zappalà, L. & Grissa-Lebdi, K. (2018c): Effectiveness of different release rates of *Trichogramma cacoeciae* (Hymenoptera: Trichogrammatidae) against *Tuta absoluta* (Lepidoptera: Gelechiidae) in protected and open field tomato crops in Tunisia. – Biocontrol Science and Technology, doi: 10.1080/09583157.2018.1542485
- Cherif, A., Attia-Barhoumi, S., Mansour, R., Zappalà, L. & Grissa-Lebdi, K. (2018d): Elucidating key biological parameters of *Tuta absoluta* on different host plants and under various temperature and relative humidity conditions. – Entomologia Generalis, doi:10.1127/entomologia/2018/0685
- Chermiti, B. & Abbes, K. (2012): Comparison of pheromone lures used in mass trapping to control the tomato leafminer *Tuta absoluta* (Meyrick, 1917) in industrial tomato crops in Kairouan (Tunisia). – EPPO Bulletin 42: 241–248.
- Chidege, M., Al-zaidi, S., Hassan, N., Julie, A., Kaaya, E. & Mrogoro, S. (2016): First record of tomato leafminer *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Tanzania. – Agriculture and Food Security 5: 17.
- Chidege, M., Abel, J., Afonso, Z., Tonini, M. & Fernandez, B. (2017): Tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) detected in Namibe Province Angola. – Journal of Applied Life Sciences International 12: 1–5.
- Cocco, A., Deliperi, S. & Delrio, G. (2013): Control of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in greenhouse tomato crops using the mating disruption technique. – Journal of Applied Entomology 137: 16–28.
- COLEACP – The Europe-Africa-Caribbean-Pacific Liaison Committee (2014): Rapport annuel du COLEACP 2014. – <https://www.coleacp.org/>, accessed on 1 August 2018.
- COLEACP – The Europe-Africa-Caribbean-Pacific Liaison Committee (2015): Rapport annuel du COLEACP 2015. – <https://www.coleacp.org/>, accessed on 1 August 2018.
- DAFF – Department of Agriculture, Forestry and Fisheries of the Republic of South Africa (2016): Media Release, Pest Alert – Detection of *Tuta absoluta* (tomato leaf miner) for the first time in the Eastern Parts of Mpumalanga Province, South Africa. – Department of Agriculture, Forestry and Fisheries, Pretoria, South Africa, <http://www.daff.gov.za.>, accessed on 27 July 2018.
- DAFF – Department of Agriculture, Forestry and Fisheries of the Republic of South Africa (2017): Media Release, Pest Alert – The update on the status of *Tuta absoluta* (tomato leaf miner) in South Africa. – Department of Agriculture, Forestry and Fisheries, Pretoria, South Africa. <http://www.daff.gov.za>, accessed on 27 July 2018.
- DAFF (2018): Abstract of Agricultural Statistics. Department of Agriculture, Forestry and Fisheries, Pretoria, South Africa. <http://www.daff.gov.za>, accessed on 27 July 2018.
- Daane, K.M., Wang, X.G., Biondi, A., Miller, B., Miller, J.C., Riedel, H. et al. (2016): First exploration of parasitoids of *Drosophila suzukii* in South Korea as potential classical biological agents. – Journal of Pest Science 89: 823–835.
- Desneux, N., Decourtye, A. & Delpuech, J.M. (2007): The sublethal effects of pesticides on beneficial arthropods. – Annual Review of Entomology 52: 81–106.

- Desneux, N., Wajnberg, E., Wyckhuys, K.A.G., Burgio, G., Arpaia, S. et al. (2010): Biological invasion of European tomato crops by *Tuta absoluta*: ecology, geographic expansion and prospects for biological control. – *Journal of Pest Science* 83: 197–215.
- Desneux, N., Luna, M.G., Guillemaud, T. & Urbaneja, A. (2011): The invasive South American tomato pinworm, *Tuta absoluta*, continues to spread in Afro-Eurasia and beyond: the new threat to tomato world production. – *Journal of Pest Science* 84: 403–408.
- Diatte, M., Brévault, T., Sylla, S., Tendeng, E., Sall-Sy, D. & Diarra, K. (2018): Arthropod pest complex and associated damage in field-grown tomato in Senegal. – *International Journal of Tropical Insect Science* 38: 243–253.
- Drouai, H., Mimeche, F., Zedam, A., Mimeche, H., Belhamra, M. & Biche, M. (2016): New floristic records of *Tuta absoluta* Meyrick 1917, in Zibans's Oasis (Biskra Algeria). – *Journal of entomology and zoology studies* 4: 130–132.
- El-Aassar, M.R., Soliman, M.H.A. & Abd Elaal, A.A. (2015): Efficiency of sex pheromone traps and some bio and chemical insecticides against tomato borer larvae, *Tuta absoluta* (Meyrick) and estimate the damages of leaves and fruit tomato plant. – *Annals of Agricultural Sciences* 60: 153–156.
- Elaini, R. (2011): Les aspects pratiques de la lutte contre *Tuta absoluta*: Cas de la région du Souss (Maroc). Control Strategies: Integrated Pest Management (IPM). – EPPO/IOBC/FAO/NEPPO Joint International Symposium on management of *Tuta absoluta* (tomato borer, Lepidoptera: Gelechiidae). – Agadir, Morocco, p 44.
- El-Arnaouty, S.A., Pizzol, J., Galal, H.H., Kortam, M.N., Afifi, A.I., Beyssat, V. et al. (2014): Assessment of Two Trichogramma Species for the Control of *Tuta absoluta* in North African Tomato Greenhouses. – *African Entomology* 22: 801–809.
- El-Lissy, O. (2014) Federal order for U.S. imports of host materials of tomato leaf miner (*Tuta absoluta*), July 23, 2014, https://www.aphis.usda.gov/import_export/plants/plant_imports/federal_order/downloads/2014/DA-2014-33.pdf, accessed on 13 July 2018.
- Eman, A.M.M., El-Refai, S.A. & Rania, S.A.R. (2016): Survey of parasitoids and predators of tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt. – *Arab Universities Journal of Agricultural Sciences* 24: 547–553.
- EPPO – European Plant Protection Organization (2008) EPPO Global Database. – <https://gd.eppo.int/>, accessed on 13 July 2018.
- EPPO – European Plant Protection Organization (2009): First report of *Tuta absoluta* in Tunisia, 2009. – EPPO Global Database, <https://archives.eppo.int/EPPOreporting/2009/Rse-0903.pdf>, accessed on 13 July 2018.
- EPPO – European Plant Protection Organization (2011): *Tuta absoluta* continues to spread around the Mediterranean Basin (Bahrain, Kuwait, Egypt, Jordan, Lebanon, Libya, Saudi Arabia and Sudan), 2011. – EPPO Global Database, <https://archives.eppo.int/EPPOreporting/2011/Rse-1104.pdf>, accessed on 20 July 2018.
- EPPO – European Plant Protection Organization (2018): *Tuta absoluta* – Distribution details in Namibia. – EPPO Global Database, <https://gd.eppo.int/taxon/GNORAB/distribution/NA>, accessed on 13 July 2018.
- Essoung, F.R.E., Mba'ning, B.M., Mohamed, S.A., Lwande, W., Ngouela, S.A., Tsamo, E. et al. (2017): Welwitschianalol A and B, two cyclohexene derivatives and other insecticidal constituents of *Caesalpinia welwitschiana* (Oliv.) Brenan. – *Phytochemistry Letters* 22: 81–86.
- Ettaib, R., Belkadhi, M.S. & Ben Belgacem, A. (2016): Study of the biotic potential of indigenous predator *Nesidiocoris tenuis* on *Tuta absoluta* pest of geothermal culture in the south of Tunisia. – *Journal of entomology and zoology studies* 4: 692–695.
- FAO – Food and Agriculture Organization of the United Nations (2015): Quarterly Early Warning Bulletin for Food and Agriculture. – No. 16, July–September 2015, <http://www.fao.org/emergencies/resources/documents/resources-detail/en/c/303378/>, accessed on 27 July 2018.
- FAO – Food and Agriculture Organization of the United Nations (2018): Food Chain Crisis, Early Warning Bulletin – Forecasting threats to the food chain affecting food security in countries and regions. – No. 28, July–September 2018, 62 p., <http://www.fao.org/3/ca0354en/CA0354EN.PDF>, accessed on 13 July 2018.

- FAOSTAT (2016): Tomato production data in 2016. FAO, Rome, Italy. – <http://www.fao.org/faostat/en/#home>
- Goda, N.F., El-Heneidy, A.H., Djelouah, K. & Hassan, N. (2015): Integrated pest management of the tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in tomato fields in Egypt. – Egyptian Journal of Biological Pest control 25: 655–661.
- Goftishu, M., Seid, A. & Dechassa, N. (2014): Occurrence and population dynamics of tomato leaf miner (*Tuta absoluta* (Meyrick), Lepidoptera: Gelechiidae) in Eastern Ethiopia. – East African Journal of Sciences 8: 59–64.
- Grisa-Lebdi, K., Skander, M., Mhafdh, M. & Belhadj, R. (2011): Lutte intégrée contre la mineuse de la tomate, *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) en Tunisie. – Faunistic Entomology 63: 125–132.
- Guedes, R.N.C. & Picanço, M.C. (2012): The tomato borer *Tuta absoluta* in South America: pest status, management and insecticide resistance. – EPPO Bulletin 42: 211–216.
- Guenauoui, Y. (2008): Nouveau ravageur de la tomate en Algérie: Première observation de *Tuta absoluta*, mineuse de la tomate invasive, dans la région de Mostaganem, au printemps 2008. – Phytoma-La Défense des Végétaux 617: 18–19.
- Guenauoui, Y., Bensaad, R. & Ouezzani, K. (2011a): Importance of native polyphagous predators able to prey on *Tuta absoluta* Meyrick (Lep: Gelechiidae) on tomato crops. – EPPO/IOBC/FAO/NEPPO Joint International Symposium on management of *Tuta absoluta*, Agadir, Morocco, p. 38.
- Guenauoui, Y., Bensaad, R., Ouezzani, K. & Vercher, R. (2011b): Emerging opportunities to use native entomophagous against *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) infesting tomato in unheated greenhouse in Northwestern Algeria. Between benefits and risks. – 9ème Conférence Internationale sur les Ravageurs en Agriculture, SupAgro, Montpellier, France, pp. 324–334.
- Guimapi, R.Y., Mohamed, S.A., Okeyo, G.O., Ndjomatchoua, F.T., Ekesi, S. & Tonnang, H.E. (2016): Modeling the risk of invasion and spread of *Tuta absoluta* in Africa. – Ecological Complexity 28: 77–93.
- Haddi, K., Berger, M., Bielza, P., Cifuentes, D., Field, L.M., Gorman, K., Rapisarda, C., Williamson, M.S. & Bass, C. (2012): Identification of mutations associated with pyrethroid resistance in the voltage-gated sodium channel of the tomato leaf miner (*Tuta absoluta*). – Insect Biochemistry and Molecular Biology 42: 506–513.
- Haddi, K., Berger, M., Bielza, P., Rapisarda, C., Williamson, M.S., Moores, G. & Bass, C. (2017): Mutation in the ace-1 gene of the tomato leaf miner (*Tuta absoluta*) associated with organophosphates resistance. – Journal of Applied Entomology 141: 612–619.
- Hafsi, A., Abbes, K., Chermiti, B. & Nasraoui, B. (2012): Response of the tomato miner *Tuta absoluta* (Lepidoptera: Gelechiidae) to thirteen insecticides in semi-natural conditions in Tunisia. – EPPO Bulletin 42: 312–316.
- Han, P., Zhang, Y.N., Lu, Z.Z., Wang, S., Biondi, A. & Desneux, N. (2018): Are we ready for the invasion of *Tuta absoluta*? Key questions remain to be answered for elaborating an Integrated Pest Management packages in Xinjiang province, China. – Entomologia Generalis, 38 (2): doi:10.1127/entomologia/2018/0739
- Han, P., Bayram, Y., Shaltiel-Harpaz, L., Sohrabi, F., Saji, A. et al. (2019): *Tuta absoluta* continues to disperse in Asia: damage, ongoing management and future challenges. – Journal of Pest Science, DOI: 10.1007/s10340-018-1062-1
- Hardwick, S., Ferguson, C.M., McNeill, M.R. & Phillips, C.B. (2016): Using mass-emergence devices to introduce an insect biocontrol agent to a new region and assist its dispersal. – Journal of Pest Sciences 89: 965–976.
- Heimpel, G.E. & Cock, M.J. (2018): Shifting paradigms in the history of classical biological control. – BioControl 63: 27–37.
- Idriss, G.E.A., Mohamed, S.A., Khamis, F., Du Plessis, H. & Ekesi, S. (2018): Biology and performance of two indigenous larval parasitoids on *Tuta absoluta* (Lepidoptera: Gelechiidae) in Sudan. – Biocontrol Science and Technology 28: 614–628.
- IPPC – International Plant Protection Convention (2014): New pest in Kenya: Preliminary surveillance report on *Tuta absoluta* – preliminary report by Kenya. – IPPC official pest report, <https://>

- www.ippc.int/es/countries/kenya/pestreports/2014/06/new-pest-in-kenya-preliminary-surveillance-report-on-tuta-absoluta-preliminary-report-by-kenya/, accessed on 3 August 2018.
- IPPC – International Plant Protection Convention (2017a): Report on tomato leafminer (*Tuta absoluta*). IPPC official pest report, <https://www.ippc.int/en/countries/ghana/pestreports/2017/07/report-on-tomato-leaf-miner-tuta-absoluta/>, accessed on 13 July 2018.
- IPPC – International Plant Protection Convention (2017b): Occurrence of tomato leaf miner (*Tuta absoluta*) in Mozambique. IPPC official pest report, <https://www.ippc.int/en/countries/Mozambique/pestreports/2017/01/occurrence-of-tomato-leaf-miner-tuta-absoluta-in-mozambique/>, accessed on 27 July 2018.
- IPPC – International Plant Protection Convention (2017c): Occurrence of South American tomato leaf miner (*Tuta absoluta*) in Malawi. – <https://www.ippc.int/en/countries/malawi/eventreporting/2017/05/occurrence-of-south-american-tomato-leaf-miner-tuta-absoluta-in-malawi-1/>, accessed on 27 July 2018
- Jacobs, D.H. (1985): Heteroptera. – In: Scholtz, C.H. & Holm, E. (eds): Insects of Southern Africa. – Butterworth Publishers, Durban, pp 112–148.
- Khidr, A.A., Gaffar, S.A., Nada, M.S., Taman, A.A. & Salem, F.A. (2013): New approaches for controlling tomato leafminer, *Tuta absoluta* (Meyrick) in tomato fields in Egypt. – Egyptian journal of agricultural research 91: 335–348.
- Konopka, J.K., Haye, T. Gariepy, T.D. & McNeil, J.N. (2017): Possible coexistence of native and exotic parasitoids and their impact on control of *Halyomorpha halys*. – Journal of Pest Sciences 90: 1119–1125.
- Kortam, M.N., El-Arnaouty, S.A., Afifi, A.I. & Heikal, I.H. (2014): Efficacy of Different Biological Methods for Controlling the Tomato Leaf Miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) on Tomato in Greenhouse in Egypt. – Egyptian Journal of Biological Pest Control 24: 523–528.
- Lietti, M.M., Botto, E. & Alzogaray, R.A. (2005): Insecticide resistance in argentine populations of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). – Neotropical Entomology 34: 113–119.
- Luangala, M.S.A., Msiska, K.K., Chomba, M.D., Mudenda, M. & Mukuwa, P.S.C. (2016): First report of *Tuta absoluta* (tomato leafminer) in Zambia. – Plant Quarantine and Phytosanitary Service, Zambia Agriculture Research Institute, Chilanga, Lusaka, 12 pp.
- Machekano, H., Mutamiswa, R. & Nyamukondiwa, C. (2018): Evidence of rapid spread and establishment of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in semi-arid Botswana. – Agriculture and Food Security 7: 48.
- Mahmoud, M.E.E. (2013): Natural enemies of *Tuta absoluta* in Kassala State, Sudan. – Paper presented at the workshop on *Tuta absoluta*: Meeting the challenge of the tomato leafminer, Addis Ababa, Ethiopia, https://www.youtube.com/watch?time_continue=134&v=il-A6EM78pY, accessed on 25 July 2018.
- Martin, T., Palix, R., Kamal, A., Deletre, E., Bonafos, R., Simon, S. & Ngouajio, M. (2013): A repellent net as a new technology to protect cabbage crops. – Journal of Economic Entomology 106:1699–1706.
- Martins, J.C., Picanço, M.C., Bacci, L., Guedes, R.N.C., Santana, P.A., Ferreira, D.O. & Chediak, M. (2016): Life table determination of thermal requirements of the tomato borer *Tuta absoluta*. – Journal of Pest Sciences 89: 897–908.
- Modiselle, S., Mosiane, S. & Nofemela, R. (2017): A serious insect pest of tomato, *Tuta absoluta*, has invaded South Africa. – Economic News 61, Agricultural Research Council, South Africa, www.arc.agric.za.
- Mohamed, S.H. (2011): Use of some eco-biological aspects and statistical analyses in determining the number of generations of *Tuta absoluta* on tomato in Upper Egypt. – EPPO/IOBC/FAO/NEPPO Joint International Symposium on management of *Tuta absoluta*, (tomato borer), Agadir, Morocco, p 68.
- Mohamed, E.S.I., Mohamed, M.E. & Gamiel, S.A. (2012): First record of the tomato leafminer, *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae) in Sudan. – EPPO Bulletin 42: 325–327.

- Mohamed, E.S.I., Mahmoud, M.E.E., Elhaj, M.A.M., Mohamed, S.A. & Ekesi, S. (2015): Host plants record for tomato leaf miner *Tuta absoluta* (Meyrick) in Sudan. – EPPO Bulletin 45: 108–111.
- Mommaerts, V., Sterk, G. & Smagge, G. (2006): Hazards and uptake of chitin synthesis inhibitors in bumblebees *Bombus terrestris*. – Pest Management Science 62: 752–758.
- Moussa, S., Sharma, A., Baiomy, F. & El-Adl, F.E. (2013): The status of tomato leafminer *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt and potential effective pesticides. – Academic Journal of Entomology 6: 110–115.
- Mutamiswa, R., Machekano, H. & Nyamukondiwa, C. (2017): First report of tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Botswana. – Agriculture and Food Security 6: 49.
- Nilahyane, A., Bouharroud, R., Hormatallah, A. & Ait Taadaouit, N. (2012): Larvicidal effect of plant extracts on *Tuta absoluta* (Lepidoptera, Gelechiidae). – IOBC/WPRS Bulletin 80: 305–310.
- ONSSA (2010): Office national de Sécurité Sanitaire des Produits. Guide pratique pour la gestion phytosanitaire de la mineuse de la tomate. <http://www.onssa.gov.ma/fr/images/pdf/2menu>, accessed on 30 May 2018.
- Oke, O.A., Kolawole, R.O., Ogunremi, O.A., Akinsola, O.A. & Awe, S.A. (2016): Detection of *apanteles* spp (Hymenoptera: Braconidae) larval parasitoid of tomato leafminer *Tuta absoluta* (Lepidoptera: Gelechiidae) on greenhouse tomato in Abeokuta, Ogun state, Nigeria. – 25th International Congress of Entomology – Program Book, Orlando, Florida, USA, p 318.
- Oke, O.A., Kolawole, R.O., Akinsola, O.A., Olajide-Taiwo, L.O., Oladigbolu, A.A. & Ajayi, E.O. (2017a): The first effort at adopting integrated pest management (IPM) to contain the infestation of *Tuta absoluta* in Nigeria. – In: Niassy, S., Ekesi, S., Migiyo, L. & Otieno, W. (eds): Sustainable Management of Invasive Pests in Africa, Springer (in press).
- Oke, O.A., Oladigbolu, A.A. & Hamisu, H.S. (2017b): The first evaluation of resistance of different insecticides on *Tuta absoluta* populations from the major tomato growing states of Nigeria. – 22nd Meeting and conference of the African Association of Insect Scientists (AAIS 2017) – Abstract Book, Wad Medani, Sudan, p 91.
- Oke, O.A. & Oladigbolu, A.A. (2018a): Natural enemies associated with *Tuta absoluta* (Meyrick) on tomato in Nigeria. – Submitted
- Oke, O.A. & Oladigbolu, A.A. (2018b): Comparative efficiency of NIHORT-Tuta trap tray against some synthetic insecticides for control of *Tuta absoluta* (Meyrick) on tomato. – Submitted
- Oke, O.A. & Oladigbolu, A.A. (2018c): Effect of integrated pest management for the control of *Tuta absoluta* (Meyrick) on tomato in the field. – Submitted
- Ouardi, K. (2011): Stratégie nationale de lutte contre la mineuse de la tomate *Tuta absoluta* Meyrick. – Power point oral presentation, Symposium international conjoint OEPP/OILB/FAO/NEPPO sur la gestion de la mineuse de la tomate *Tuta absoluta*, Agadir, Morocco.
- Ouardi, K., Chouibani, M., Rahel, M.A. & El Akel, M. (2012): Stratégie Nationale de lutte contre la mineuse de la tomate *Tuta absoluta* Meyrick. – EPPO Bulletin 42: 281–290.
- Peris, N.W., Muturi, J.J., Mark, O., Esther, A. & Jonsson, M. (2018): Tomato Leaf miner (*Tuta absoluta*) (Meyrick 1917) (Lepidoptera: Gelechiidae) prevalence and farmer management practices in Kirinyanga County, Kenya. – Journal of Entomology and Nematology 10: 43–49.
- Pfeiffer, D.G., Muniappan, R., Sall, D., Diatta, P., Diongue, A. & Dieng, E.O. (2013): First record of *Tuta absoluta* (Lepidoptera: Gelechiidae) in Senegal. – Florida Entomologist 96: 661–662.
- Pimentel, D., Lach, L., Zuniga, R. & Morrison, D. (2000): Environmental and economic costs of nonindigenous species in the United States. – BioScience 50: 53–65.
- Pimentel, D., Zuniga, R. & Morrison, D. (2005): Update on the environmental and economic costs associated with alien-invasive species in the United States. – Ecological Economics 52: 273–288.
- Retta, A.N. & Berhe, D.H. (2015): Tomato leaf miner *Tuta absoluta* (Meyrick), a devastating pest of tomatoes in the highlands of Northern Ethiopia: A call for attention and action. – Research Journal of Agricultural and Environmental Management 4: 264–269.

- Reyes, M., Rocha, K., Alarcón, L., Siegwart, M. & Sauphanor, B. (2012): Metabolic mechanisms involved in the resistance of field populations of *Tuta absoluta* (Meyrick) (Lep., Gelechiidae) to spinosad. – *Pesticide Biochemistry and Physiology* 102: 45–50.
- Rizk, A.M. (2016): Effectiveness of different bio-techniques for controlling the pin worm, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). – *Egyptian Journal of Biological Pest control* 26: 797–802.
- Roditakis, E., Skarmoutsou, C. & Staurakaki, M. (2013a): Toxicity of insecticides to populations of tomato borer *Tuta absoluta* (Meyrick) from Greece. – *Pest Management Science* 69: 834–40.
- Roditakis, E., Skarmoutsou, C., Staurakaki, M., del Rosario Martínez Aguirre, M., García Vidal, L. et al. (2013b): Determination of baseline susceptibility of European populations of *Tuta absoluta* (Meyrick) to indoxacarb and chlorantraniliprole using a novel dip bioassay method. – *Pest Management Science* 69: 217–227.
- Roditakis, E., Vasakis, E., Grispou, M., Stavrakaki, M., Nauen, R. et al. (2015): First report of *Tuta absoluta* resistance to diamide insecticides. – *Journal of Pest Science* 88: 9–16.
- Roditakis, E., Vasakis, E., García-Vidal, L., del Rosario Martínez-Aguirre, M., Rison, J.L. et al. (2018): A four-year survey on insecticide resistance and likelihood of chemical control failure for tomato leaf miner *Tuta absoluta* in the European/Asian region. – *Journal of Pest Science* 91: 421–435.
- Sala, O.E., Chapin III, F.S., Armesto, J.J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-Sanwald, E., Huenneke, L.F., Jackson, R.B., Kinzig, A., Leemans, R., Lodge, D.M., Mooney, H.A., Oesterheld, M., Poff, N.L., Sykes, M.T., Walker, B.H., Walker, M. & Wall, D.H. (2000): Global biodiversity scenarios for the year 2100. – *Science* 287: 1770–1774.
- Salama, H.S.A., Ismail, I.A., Fouda, M., Ebadah, I. & Shehata, I. (2015): Some Ecological and Behavioral Aspects of the Tomato Leaf Miner *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). – *Journal of Balkan Ecology* 7: 35–44.
- Salas Gervassio, N.G., Aquino, D., Vallina, C., Biondi, A. & Luna, M.G. (2019): A re-examination of *Tuta absoluta* parasitoids in South America for optimized biological control. – *Journal of Pest Science*.
- Salazar, E.R. & Araya, J.E. (1997): Detección de resistencia a insecticidas en la polilla del tomate (Detection of insecticide resistance in the tomato moth). – *Simiente* 67: 8–22.
- Salazar, E.R. & Araya, J.E. (2001): Tomato moth, *Tuta absoluta* (Meyrick) response to insecticides in Arica, Chile. – *Agricultura Técnica* 61: 429–435.
- Salehi, Z., Yarahmadi, F., Rasekh, A. & Sohani, N.Z. (2016): Functional responses of *Orius albidipennis* Reuter (Hemiptera, Anthocoridae) to *Tuta absoluta* Meyrick (Lepidoptera, Gelechiidae) on two tomato cultivars with different leaf morphological characteristics. – *Entomologia Generalis* 36 (2): 127–136.
- Sanda, N.B., Sunusi, M., Hamisu, H.S., Wudil, B.S., Sule, H. & Abdullahi, A.M. (2018): Biological invasion of tomato leafminer, *Tuta absoluta* (Meyrick) in Nigeria: problems and management strategies optimization: A Review. – *Asian Journal of Agricultural and Horticultural Research* 1: 1–14.
- Sarto, M.C.L.D., Oliveira, E.E., Guedes, R.N.C. & Campos, L.A.O. (2014): Differential insecticide susceptibility of the Neotropical stingless bee *Melipona quadrifasciata* and the honey bee *Apis mellifera*. – *Apidologie* 45: 626–636.
- Shiberu, T. & Getu, E. (2018): Evaluation of bio-pesticides on integrated management of tomato leafminer, *Tuta absoluta* (Meyrick) (Gelechiidae: Lepidoptera) on tomato crops in Western Shewa of Central Ethiopia. – *Entomology, Ornithology and Herpetology* 7: 1.
- Silva, T.B., Silva, W.M., Campos, M.R., Silva, J.E., Ribeiro, L.M., Siqueira & H.A.A. (2016): Susceptibility levels of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) to minor classes of insecticides in Brazil. – *Crop Protection* 79: 80–86
- Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J., Courchamp, F., Galil, B., Garcia-Berthou, E., Pascal, M., Pyšek, P., Sousa, R., Tabacchi, E. & Vilà M (2013): Impacts of biological invasions: what's what and the way forward. – *Trends in Ecology and Evolution* 28: 58–66.

- Siqueira, H.A., Guedes, R.N. & Picanço, M.C. (2000): Cartap resistance and synergism in populations of *Tuta absoluta* (Lep., Gelechiidae). – *Journal of Applied Entomology* 124: 233–238.
- Sohrabi, F., Nooryazdan, H., Gharati, B. & Saeidi, Z. (2016): Evaluation of ten tomato cultivars for resistance against tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) under field infestation conditions. – *Entomologia Generalis* 36 (2):163–175.
- Son, D., Bonzi, S., Somda, I., Bawin, T., Boukraa, S., Verheggen, F., Francis, F., Legrève, A. & Schiffers, B. (2017): First record of *Tuta absoluta* (Meyrick, 1917) (Lepidoptera: Gelechiidae) in Burkina Faso. – *African Entomology* 25: 259–263.
- Son, D., Zerbo, F.K., Bonzi, S., Legreve, A., Somda, I. & Schiffers, B. (2018): Assessment of Tomato (*Solanum lycopersicum* L.) Producers' Exposure Level to Pesticides, in Kouka and Toussiana (Burkina Faso). – *International Journal of Environmental Research and Public Health* 15 (2), doi: 10.3390/ijerph15020204
- Sylla, S., Brévault, T., Streito, J.C. & Diarra, K. (2016): First Record of *Nesidiocoris tenuis* (Reuter) (Heteroptera: Miridae), as a predator of the tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae), in Senegal. – *Egyptian Journal of Biological Pest Control* 26: 851–853.
- Sylla, S., Brévault, T., Bal, A.B., Chailleux, A., Diatte, M., Desneux, N. & Diarra, K. (2017): Rapid spread of the tomato leafminer, *Tuta absoluta* (Lepidoptera: Gelechiidae), an invasive pest in Sub-Saharan Africa. – *Entomologia Generalis* 36 (3): 269–283.
- Sylla, S., Seydi, O., Diarra, K. & Brévault, T. (2018): Seasonal decline of the tomato leafminer, *Tuta absoluta*, in the shifting landscape of a vegetable-growing area. – *Entomologia Experimentalis et Applicata*, doi:10.1111/eea.12722
- Taha, A.M., Afsah, A.F.E. & Fargalla, F.H. (2013): Evaluation of the effect of integrated control of tomato leafminer *Tuta absoluta* with sex pheromone and insecticides. – *Natural Sciences* 11: 26–29.
- Tena, A., García-Bellón, J. & Urbaneja, A. (2017): Native and naturalized mealybug parasitoids fail to control the new citrus mealybug pest *Deltoecoccus aberiae*. – *Journal of Pest Sciences* 90: 659–667.
- Tonnang, H.E., Mohamed, S.F., Khamis, F. & Ekesi, S. (2015): Identification and risk assessment for worldwide invasion and spread of *Tuta absoluta* with a focus on Sub-Saharan Africa: implications for phytosanitary measures and management. – *PloS One* 10: e0135283.
- Tropea Garzia, G., Siscaro, G., Biondi, A. & Zappalà, L. (2012): *Tuta absoluta*, a South American pest of tomato now in the EPPO region: Biology, distribution and damage. – *EPPO Bulletin* 42: 205–210.
- USDA (2011): New Pest Response Guidelines – Tomato Leafminer (*Tuta absoluta*), United States Department of Agriculture, Animal and Plant Health Inspection Service, Cooperating State Departments of Agriculture, 176 pp., https://www.aphis.usda.gov/import_export/plants/manuals/emergency/downloads/Tuta-absoluta.pdf, accessed on 13 July 2018.
- Vacas, S., Alfaro, C., Primo, J. & Navarro-Llopis, V. (2011): Studies on the development of a mating disruption system to control the tomato leafminer, *Tuta absoluta* Povolny (Lepidoptera: Gelechiidae). – *Pest Management Science* 67: 1473–80.
- Valencia, L. & Peñaloza, J. (1990): Control Biológico de las palomillas de la papa. Periódico rural. – “El Boyacense”, Tunja, Colombia, Ministerio de Agricultura, ICA: 4 p.
- Vallejo, F.A. (1999): Mejoramiento genético y producción de tomate en Colombia. – Universidad Nacional de Colombia, Sede Palmira, Cali (Colombia), pp. 17–22, 35, 100, 101, 160–167.
- Visser, D., Uys, V.M., Nieuwenhuis, R.J. & Pieterse, W. (2017): First records of the tomato leaf miner *Tuta absoluta* (Meyrick, 1917) (Lepidoptera: Gelechiidae) in South Africa. – *BioInvasions Records* 6: 301–305.
- Wanumen, A.C., Carvalho, G.A., Medina, P., Viñuela, E. & Adán, A. (2016): Residual acute toxicity of some modern insecticides toward two mirid predators of tomato pests. – *Journal of Economic Entomology* 109: 1079–85.
- Xian, X.Q., Han, P., Wang, S., Zhang, G.F., Liu, W.X., Desneux, N. & Wan, F. (2017): The potential invasion risk and preventive measures against the tomato leafminer *Tuta absoluta* in China. – *Entomologia Generalis* 36 (4): 319–333.

- Zappalà, L., Biondi, A., Alma, A., Al-Jboory, I.J., Arno, J. et al. (2013): Natural enemies of the South American moth, *Tuta absoluta*, in Europe, North Africa and Middle-East, and their potential use in pest control strategies. – *Journal of Pest Sciences* 86: 635–647.
- Zouba, A., Chermiti, B., Chraïet, R. & Mahjoubi, K. (2013a): Effect of Two Indigenous *Trichogramma* Species on the Infestation Level by Tomato Miner *Tuta absoluta* in Tomato Greenhouses in the South-West of Tunisia. – *Tunisian Journal of Plant Protection* 8: 87–106.
- Zouba, A., Chermiti, B., Kadri, K. & Fattouch, S. (2013b): Molecular characterization of *Trichogramma bourarachae* strains (Hymenoptera: Trichogrammatidae) from open field tomato crops in the South West of Tunisia. – *Biomirror* 4: 5–11.

Manuscript received: 30 July 2018

Accepted: 1 October 2018